

A Variation of the Present Star Formation Activity of Spiral Galaxies

Akihiko TOMITA, Yoshio TOMITA, and Mamoru SAITŌ

*Department of Astronomy, Faculty of Science, Kyoto University,
Sakyo-ku, Kyoto 606-01*

E-mail (AT) atomita@kusastro.kyoto-u.ac.jp

(Received 1995 November 7; accepted 1996 January 5)

To be appeared in PASJ, Vol. 48, No. 2 (April 1996 issue)

Abstract

The star formation rate in spiral galaxies is considered to be decreasing continuously with time in a time scale of 10^9 yr. The present star formation activity, on the other hand, occurs in molecular clouds with a time scale of 10^7 yr, and shows various degrees among galaxies. We make a new data set of 1681 nearby spiral galaxies from available databases and study the statistics of the present star formation activity. We analyze far-infrared and optical B -band surface brightnesses of the H II regions and the non-H II regions in M 31 and show that a far-infrared-to-optical B -band flux ratio, $f_{\text{FIR}}/f_{\text{B}}$, is a useful indicator of the present star formation activity of spiral galaxies. For the sample galaxies, we make the distribution diagram of $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ for each morphological type. The distribution of $f_{\text{FIR}}/f_{\text{B}}$ has a dispersion of one to two orders of magnitude even within the same morphological type of galaxies, implying that the star formation activities of spiral galaxies changes discontinuously in a short time scale. Analyzing the $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ correlation, we suggest that the most active star formation in galaxies does not continue longer than 10^8 yr. We also construct a universal distribution histogram of $\log (f_{\text{FIR}}/f_{\text{B}})$ for each morphological type. The earlier-type spirals tend to show larger variation of the present star formation activity. We discuss the correlation between the present star formation activity and the structures and environments of galaxies. We suggest that the short-term variation occurs primarily due to internal processes which may change with the morphological type of galaxies.

Key words: Galaxies: star formation — Galaxies: evolution — Galaxies: star formation indicators — Galaxies: morphological types

1 Introduction

The star formation rate (SFR) in spiral galaxies with a time scale of 10^9 yr seems to be continuously decreasing with time. The late-type spirals have reduced the SFR with a time scale comparable to the Hubble time (e.g., Searle et al. 1973; Kennicutt 1983; Gallagher et al. 1984), while the early-type spirals have a low preset SFR compared with the past SFR and the decaying time scale is much shorter than the Hubble time scale (e.g., Kennicutt 1983, Sandage 1986). Photometric evolution models under such a view of the star formation histories reproduced the present colors of galaxies along the Hubble sequence (e.g., Bruzual 1983; Arimoto, Yoshii 1986; Guiderdoni, Rocca-Volmerange 1987). Stars, especially massive stars, mostly form in giant molecular clouds (GMC). The time scale of the star formation activity in a GMC is of order of 10^7 yr (e.g., Shu et al. 1987), and the dynamical time scale of a galaxy is $\sim 10^8$ yr; these are much shorter than the time scale of Gyr. In the galaxy scale, some galaxies show transient starburst phenomenon and there are post-starburst galaxies (e.g., Walker et al. 1988; Beckman et al. 1991). These suggest that the active star formation occurs discontinuously in a short time scale. This has been also suggested by statistical studies of far-infrared (FIR)-to-optical B -band flux ratios of galaxies (e.g., de Jong et al. 1984; Soifer et al. 1987; Bothun et al. 1989).

It is also a controversial question what kind of global parameters of galaxies are closely related to the present star formation activity. Roberts et al. (1979) and Kenney et al. (1992) argued that the bar structure triggers a nuclear starburst. However, Moss and Whittle (1993) showed from analysis of the $H\alpha$ equivalent width that the existence of bar is not correlated with the present star formation activity. Dressler et al. (1985) claimed that cluster galaxies have smaller star formation activity than field galaxies, while Gavazzi and Jaffe (1985) and Moss and Whittle (1993) concluded that cluster galaxies have more intense star formation activity. The galaxy-galaxy interaction is considered to be an effective triggering mechanism for starburst (e.g., Keel 1993), while Campos-Aguilar and Moles (1991) suggested that occurrence of the starburst is independent of the galaxy-

galaxy interaction.

The best way to study the star formation features in a short time scale and its connection with the global characteristics of galaxies is to make a reliable indicator of the present star formation activity and to investigate the difference of the present star formation activity among various galaxies. In this paper, we construct a new large high-quality data set of spiral galaxies using available databases. We describe the sample in section 2. In section 3 we make a far-infrared (FIR)-to-optical B -band flux ratio, $f_{\text{FIR}}/f_{\text{B}}$, for each galaxy and show a systematic distribution of the flux ratios of the sample galaxies. Using the measured FIR and B band surface brightnesses at various areas of M 31, we confirm that the $f_{\text{FIR}}/f_{\text{B}}$ is a useful measure of the present star formation activity in the galaxy scale. We present for the first time a distribution of $\log (f_{\text{FIR}}/f_{\text{B}})$ of spiral galaxies for each morphological type. The distributions of $f_{\text{FIR}}/f_{\text{B}}$ show that the present star formation activity has a great variety even within the same morphological type of galaxies and suggest that the star formation activity in a short time scale changes discontinuously. In section 4 we discuss what kind of global features of galaxies are correlated to the present star formation activity. A summary is given in section 5.

2 The Sample

We are interested in the star formation histories of spiral galaxies and select a sample of spirals using *Third Reference Catalogue of Bright Galaxies* (de Vaucouleurs et al. 1991; hereafter RC3). RC3 is the largest optical galaxy catalog describing various parameters of galaxies. We made use of *NAO/ADAC CD-ROM Catalog Volume 1* to search the RC3 data.

The B magnitudes of galaxies provide a measure of the galaxies and nearly represent the total mass (e.g., Roberts, Haynes 1994). RC3 gives an extinction-corrected B magnitude of each galaxy which is denoted as B_{T}^0 and corresponds to B magnitude as is observed at face-on. In the following analyses, we also use the data of the color $(B - V)_{\text{T}}^0$,

the morphological index T , and the radial velocity in RC3; the optical color of the galaxy is considered to be a result of the star formation history in the galaxy (e.g., Larson, Tinsley 1978; Arimoto, Yoshii 1986; Guiderdoni, Rocca-Volmerange 1987), the morphological type is one of the fundamental parameters of galaxies (e.g., Roberts, Haynes 1994), and the radial velocity is used to estimate the distance and the luminosity of the galaxy. Out of 13537 galaxies with the morphological index of $0 \leq T \leq 11$ (spirals and irregulars) cataloged in RC3, 1782 have all values of B_T^0 , $(B - V)_T^0$, and radial velocity. We select galaxies regardless of the bar structure. We also use the far-infrared (FIR) emission fluxes of galaxies measured by *IRAS* as an essential parameter of this study. Out of the 1782 galaxies, 101 lie in the non-surveyed regions of *IRAS* mission or belong to close binary systems for which we do not identify the association of the *IRAS* source. We thus finally adopt 1681 galaxies.

Table 1 shows the morphology and magnitude distribution of the sample galaxies and table 2 shows the morphology and radial velocity distribution; about 80% of the sample galaxies are brighter than 14 mag and have redshifts less than 5000 km s^{-1} . The sample fairly covers all of the morphological types. Figure 1 shows a distribution histogram of $(B - V)_T^0$ for each morphological type. There is a correlation between the $(B - V)_T^0$ color and the morphology; the later the morphological type is, the bluer the $(B - V)_T^0$ color is. The variation from S0a to Sc is larger than that among later types. The difference of the median of the $(B - V)_T^0$ color between the adjacent morphological types is comparable to the dispersion of the $(B - V)_T^0$ color in each morphological type, $\Delta (B - V)_T^0 \sim 0.3$. These color characteristics along the spiral Hubble sequence have been known (e.g., Sauvage, Thuan 1994; Roberts, Haynes 1994). Our sample also has the same trend.

3 Star Formation Activity in a Short Time Scale

3.1 An Indicator of Star Formation Activity in Spiral Galaxies

It is well known that the $H\alpha$ equivalent width is a direct indicator of the present

star formation rate (e.g., Kennicutt 1983). However, the $H\alpha$ emission suffers from the extinction by interstellar medium (ISM) and it is difficult to obtain the total $H\alpha$ flux of the galaxy (Kennicutt 1992). The FIR emission of spiral galaxies originates from both warm dust heated by massive stars and cool dust heated by general stellar field (Presson, Helou 1987; Devereux, Young 1990; Sauvage, Thuan 1992; Devereux, Scowen 1994; Devereux et al. 1994; Devereux et al. 1995). Gavazzi et al. (1991) showed that the FIR flux normalized to optical flux has a correlation with the $H\alpha$ equivalent width in two orders of magnitude. Xu (1990) suggested that radiation from dust heated by UV continuum dominates the total FIR emission of spiral galaxies. These indicate that the FIR flux is also an indicator of the present star formation rate in spiral galaxies. The FIR-to- B flux ratio has been used as an indicator for the present star formation activity in galaxies (de Jong et al. 1984; Fabbiano et al. 1988; Soifer et al. 1987; Bothun et al. 1989; Sauvage, Thuan 1994). Belfort et al. (1987) also showed from their photometric synthesis model that the FIR-to- B flux ratio responds to a recent starburst within 2×10^7 yr.

Following Soifer et al. (1987), we define the B -band flux f_B as 4400 Å times flux density [$\text{W m}^{-2} \text{Å}^{-1}$] at 4400 Å. Using an equation between the B -band flux and the magnitude system given in Allen (1973), we obtain $\log f_B [\text{W m}^{-2}] = -0.4 B_T^0 - 7.53$.

The *IRAS* survey has provided us FIR fluxes for a large, homogeneous, nearly all-sky, flux-limited sample, which are free from the heavy extinction by the ISM. The total FIR fluxes for galaxies are available using following five *IRAS* catalogs. The FIR fluxes of the most sample galaxies are taken from *IRAS Faint Source Catalog* (FSC) and *IRAS Point Source Catalog* (PSC). Since the measured aperture of *IRAS* point sources is about 2 arcmin, we underestimate the FIR fluxes for the galaxies with angular sizes larger than 2 arcmin. We refer to *IRAS Small Scale Structure Catalog* (SSSC) for the galaxies with sizes of about 2 to 8 arcmin. For the galaxies with sizes larger than 8 arcmin, we adopt the data of Rice et al. (1988). For the galaxies having 60 μm flux densities greater than 5.24 Jy, we take the data of the FIR-bright galaxies by Soifer et al. (1989).

Cross identification with optical objects was performed mainly based on the description in the above-mentioned catalogs and *Cataloged Galaxies and Quasars in the IRAS Survey* (hereafter CGQIRAS) and it was also checked using NASA/IPAC Extragalactic Database (NED). We made use of *ADC CD-ROM Selected Astronomical Catalogs Volume 1* in searching the data of PSC, FSC, SSSC, and CGQIRAS. We calculate ¹ the FIR flux from the flux densities at 60 and 100 μm , f_{60} and f_{100} , following CGQIRAS; i.e., $\log f_{\text{FIR}} (42.5 - 122.5 \mu\text{m}) [\text{W m}^{-2}] = \log (2.58 f_{60} [\text{Jy}] + f_{100} [\text{Jy}]) - 13.90$. 1150 galaxies have reliable measurements in both f_{60} and f_{100} . In figure 2, we obtain an empirical formula¹, $\log (f_{100}/f_{60}) = -0.23 \log (f_{60} [\text{Jy}]/f_{\text{B}} [\text{W m}^{-2}]) + 3.41$ with a correlation coefficient of 0.72, from 1150 galaxies with reliable f_{60} and f_{100} . Objects with relatively large deviations are mostly late-type spirals or irregulars; a correlation coefficient using only objects with $0 \leq T \leq 5$ is 0.80. Using the regression line, we estimate f_{100} for 156 objects with upper-limited f_{100} and reliable f_{60} , and calculate f_{FIR} of these objects. The FIR fluxes estimated by this method have an error of 4% in $\log (f_{100}/f_{60})$, which is negligible in the following analyses. An upper limit of $\log f_{\text{FIR}} = -13.82 [\text{W m}^{-2}]$ is given for 375 sample galaxies without *IRAS* detection adopting the detection limits of *IRAS* survey of $f_{60} = 0.2 \text{ Jy}$ and $f_{100} = 0.7 \text{ Jy}$ (FSC).

3.2 $f_{\text{FIR}}/f_{\text{B}}$ vs f_{100}/f_{60} Characteristics at Various Regions of M31

Figure 3 shows a $\log (f_{100}/f_{60})$ versus $\log (f_{\text{FIR}}/f_{\text{B}})$ relation for the 1150 sample galaxies which have reliable f_{60} and f_{100} values. The color temperatures corresponding to $\log (f_{100}/f_{60})$ shown in the ordinate is about $41.7 / [1 + \log (f_{100}/f_{60})]$ K for a modified black-body radiation in the form of $\nu^{1.5} B_{\nu}$, which gives about 42 K, 30 K, and 23 K for $\log (f_{100}/f_{60}) = 0, 0.4$, and 0.8 , respectively. The values of $f_{\text{FIR}}/f_{\text{B}}$ shown in the abscissa are distributed over two orders of magnitude and the galaxies with higher values of $f_{\text{FIR}}/f_{\text{B}}$ tend to have smaller f_{100}/f_{60} values, i.e., higher temperatures; the correlation

¹ The readers would notice the dimension discrepancies in these equations. These are only arithmetical formulae.

coefficient is 0.63 for the total 1150 sample and 0.72 for 916 objects with $0 \geq T \geq 5$, shown by larger circles in figure 3. The $\log(f_{100}/f_{60})$ versus $\log(f_{60}/f_B)$ relation has a somewhat tighter correlation (see section 3.1 and figure 2), because f_{60} is more insensitive for the “cirrus” component than f_{100} (e.g., Presson, Helou 1987). In figure 3, six famous spiral galaxies without strong AGN are denoted by filled squares with numbers, 1: M 31, 2: M 81, 3: M 33, 4: M 101, 5: M 51, and 6: M 83.

de Jong et al. (1984) presented that the values of the f_{FIR}/f_B of galaxies are distributed in two orders of magnitude by analyzing 165 optically bright galaxies. Soifer et al. (1987) showed that the f_{FIR}/f_B of 313 FIR-bright galaxies are independent of the B luminosity over two orders of magnitude, and suggested that the star formation activity of galaxies considerably varies in a short time scale. Bothun et al. (1989) confirmed that the values of f_{FIR}/f_B distribute over two orders of magnitude among about 3500 UGC galaxies. However, Bothun et al. (1989) also claimed that it is dangerous to take the f_{FIR}/f_B value as a direct indicator of the present star formation activity because “cirrus” component may contribute much in the FIR emission. We examine the usefulness of the f_{FIR}/f_B value as an indicator of the present star formation activity using the data of M 31.

Rice (1993) made *IRAS Nearby Galaxy High Resolution Image Atlas* (hereafter *IRAS HiRes*) of 30 galaxies for *IRAS* four bands with a 1 arcmin resolution, using an image reconstruction technique; among them there are $2^\circ \times 2^\circ$ (481×481 pixels) FITS images of M 31 ($T = 3.0$). Because of the proximity of M 31, their *IRAS* images can separate each of giant H II regions. The maps are distributed as Infrared Processing and Analysis Center (IPAC) released products. We made use of the on-line service via Automated Retrieval Mail System (ARMS) at NASA’s Data Archive Distribution Service (NDADS). One of us (Y. T.) and H. Suzuki obtained a B -band image of M 31 at Ouda Station, Kyoto University using the Wide Field CCD Camera of $f_l = 300$ mm and $F/4.5$; its field of view is $2.^\circ5 \times 1.^\circ7$ (576×384 pixels). The observation was made at October 27, 1992 and the total exposure time was 20 min. The sky subtraction was made with SPIRAL

(Hamabe, Ichikawa 1992) and foreground stars were removed by using an IRAF² task, ‘imedit’. The B -band image of M 31 is shown in figure 4 (a). Figure 4 (b) shows f_{FIR}/f_B image of M 31, made by using the B -band image and the FIR image constructed from f_{60} and f_{100} images of *IRAS HiRes*. From the f_{FIR}/f_B and f_{100}/f_{60} ratio maps, we obtain the ratios at some areas of H II regions in disk (2 to 8 kpc), non-H II regions in disk (2 to 8 kpc), bulge region (1 to 2 kpc), and nuclear region (within 0.3 kpc), where the distances in parentheses are measured from the nucleus of M 31 and the aperture radius is 0.′5. The sample locations are shown on the B -band and f_{FIR}/f_B images of figure 4.

Figure 5 shows the $\log (f_{\text{FIR}}/f_B)$ versus $\log (f_{100}/f_{60})$ relation for the various locations in M 31, where the extinction in B band was assumed to be $\Delta \log f_B = 0.4$. Both the non-H II regions in the disk and the bulge region are located at similar positions on this diagram and we simply call both of them as the non-H II regions, which correspond to the cirrus region. The plots in figure 5 are concentrated into three regions, which are the H II regions (A), the non-H II regions (B), and the central regions (C). A big star mark in figure 5 indicates the whole of M 31. The three regions and the whole M 31 are also superimposed on figure 3 which shows the $\log (f_{\text{FIR}}/f_B)$ versus $\log (f_{100}/f_{60})$ relation of the sample galaxies. In M 31, when the aperture size is increased outward the H II regions, the values of f_{100}/f_{60} and f_{FIR}/f_B move on the diagram of figure 5 from the region A to the region B. A solid line in figure 3 indicates the path of the plots generated by increasing the aperture size.

In figure 5, the nuclear region occupies different positions from the H II region and the non-H II region positions. Only 4% of the total FIR emission comes from the nuclear region in M 31 (Devereux et al. 1994), and the nuclear region has little contribution to the colors of the whole of M 31. There are a small number of galaxies among our sample whose FIR emissions are heavily contaminated by AGN. We referred to NED to search for the objects with AGN. We found that only 3.6% (34 objects) of our total sample are

² IRAF is the software developed in National Optical Astronomy Observatories, USA.

objects with AGN including possible and suspect ones. Most of AGNs are LINERs or type-2 Seyferts, i.e., weak AGNs. Even for M 81, a galaxy with LINER, the nuclear region emits only 17% of the total FIR emission (Devereux et al. 1995). Though we did not exclude the objects with AGNs from our sample, the AGN-contamination affects little on the following analyses.

We find in figure 3 that galaxies with the smallest $f_{\text{FIR}}/f_{\text{B}}$ are located around the non-H II region (B) area, and that the galaxy distribution on the diagram is along the path shown by the solid line. In figure 3, the active star-forming galaxies, M 101 (No. 4), M 51 (No. 5), and M 83 (No. 6) lie in the region A and the inactive star-forming galaxy M 31 (No. 1) lies in the region B. The order from No. 1 (M 31) to No. 6 (M 83) represents the magnitude of the star formation activities usually accepted for those galaxies (e.g., Kennicutt, Kent 1983; Kennicutt et al. 1989; Kennicutt et al. 1994). Beichman (1987) made an f_{25}/f_{60} versus f_{60}/f_{100} diagram of galaxies and showed that galaxies on this color-color diagram are located between the cold cirrus region and the hot H II region, consistent with our analysis in M 31. The spatial resolution of 1 arcmin in *IRAS HiRes* corresponds to about 200 pc on M 31, which is a little larger than typical H II region size of dozens to hundred pc. Therefore, the values of $f_{\text{FIR}}/f_{\text{B}}$ derived for the H II regions are somewhat smaller than that the true values of the H II regions and the true H II regions may move toward lower-right on the diagram of figure 5. The H II regions in M 31 do not contain early O-type stars (Devereux et al. 1994). Wood and Churchwell (1989) found that ultra-compact H II regions in the Galaxy have $\log (f_{100}/f_{60}) = 0$ to 0.5, which are similar to those of the region A of M 31. Thus, the region A of M 31 represents the position of universal H II region on the diagram. Moreover, most of the *IRAS* sources associated with low-mass young stellar objects have values of $\log (f_{100}/f_{60})$ similar to those of the region A (e.g., Beichman et al. 1986). The region A seems to represent the position of the star-forming region in general.

We thus conclude that $f_{\text{FIR}}/f_{\text{B}}$ is a useful indicator of the present star formation ac-

tivity in the whole galaxy, although the influence of the cirrus component is contained in the value of f_{FIR} . We show in appendix 1 that $f_{\text{FIR}}/f_{\text{B}}$ is not affected by the internal absorption of galaxy by checking a correlation between the distribution of $\log (f_{\text{FIR}}/f_{\text{B}})$ and apparent inclination of galaxies. It is advantageous at the present time that the indicators are available with reliable values for a large number of galaxies compared with other indicators, such as the $\text{H}\alpha$ equivalent widths. The zero-point, i.e., without the present star formation, is at $\log (f_{\text{FIR}}/f_{\text{B}}) \simeq -2$, and $f_{\text{FIR}}/f_{\text{B}}$ becomes larger with increasing of the present star formation activity and in this case, following Devereux and Young (1991) and Sauvage and Thuan (1992), the present SFR [$M_{\odot} \text{ yr}^{-1}$] $\simeq 1.4 \times 10^{-10} L_{\text{FIR}} [L_{\odot}]$. Gallagher et al. (1984) derived a relation between the B luminosity and the past-averaged SFR, as $\text{SFR} [M_{\odot} \text{ yr}^{-1}] \simeq 0.3 \times 10^{-10} L_{\text{B}} [L_{\odot}]$ under an assumption that the SFR is constant during 10^9 yr order and the upper-mass cut-off of $100 M_{\odot}$. If we use these relations, the present-to-past averaged SFR (SFR-ratio) is expressed as $\sim (1.4/0.3) (L_{\text{FIR}}/L_{\text{B}}) = (1.4/0.3) (f_{\text{FIR}}/f_{\text{B}})$; the SFR-ratio = 1, a constant SFR during 10^9 yr order, corresponds to $\log (f_{\text{FIR}}/f_{\text{B}}) \sim -0.6$ to -0.7 .

3.3 Distribution of $f_{\text{FIR}}/f_{\text{B}}$ of Galaxies

Figure 6 shows a $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ relation for each morphological index T given in RC3 from 0: S0a to 10: Im. The blue luminosity is calculated by $L_{\text{B}} = 4\pi D^2 f_{\text{B}}$, where D is the distance obtained by V_{GSR}/H_0 for the Hubble constant $H_0 = 75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ and V_{GSR} of the recession velocity [km s^{-1}] with respect to Galactic center (Galaxy Standard of Rest) taken from RC3. When V_{GSR} is smaller than 75 km s^{-1} , we give $D = 1 \text{ Mpc}$; the number of objects with $V_{\text{GSR}} < 75 \text{ km s}^{-1}$ is 19 and only 1% of the total sample. We include one object of $T = 11$ (cI) into the $T = 10$ (Im) group. Circles indicate objects with *IRAS* detection, total number is 1306, including objects without reliable f_{100} but f_{FIR} of which are estimated by the procedure mentioned in section 3.1. Crosses show the upper limits of $f_{\text{FIR}}/f_{\text{B}}$ for the objects without *IRAS* detection; the total number is 375, and the true positions are lower than the positions plotted. The

size of the symbols represents the B_T^0 magnitude of the object, as shown at the upper left in each diagram. Near the left ordinate, eight small tick marks are shown. They indicate the detection limits of f_{FIR}/f_B , due to the *IRAS* detection limits, for a given B_T^0 magnitude, $B_T^0 = 16, 15, 14, 13, 12, 11, 10$, and 9 mag, respectively, from upper to lower. For instance, an object with $B_T^0 = 11$ is detectable by *IRAS* if $\log (f_{\text{FIR}}/f_B) > -2$.

From figure 6, we confirm not only the earlier claim by others (e.g., Soifer et al. 1987) that the values of $\log (f_{\text{FIR}}/f_B)$ distribute over two orders of magnitude but also that the wide dispersion of f_{FIR}/f_B is seen even in each morphological type. The trend that the distributions of f_{FIR}/f_B show no systematic change along the morphological type is in contrast with the systematic change of such as $(B - V)_T^0$ color (see figure 1). If the SFR is constant during $\sim 10^9$ yr in a galaxy, $\log (f_{\text{FIR}}/f_B)$ would be a constant regardless of L_B . This is not the case in figure 6, indicating that the star formation activity in the galaxy scale is variable with a shorter time. We estimate in appendix 2 that the time scale of the variation is shorter than about 10^8 yr.

It is an interesting feature that f_{FIR}/f_B is independent of L_B in early-type spirals (S0a to Sc), while in late-type spirals (later than Scd, i.e., $T \geq 6$) f_{FIR}/f_B tends to decrease with decreasing L_B ; figure 7 shows $\log (f_{\text{FIR}}/f_B)$ versus $\log L_B$ diagrams for the $0 \leq T \leq 5$ group and the $6 \leq T \leq 11$ group, respectively. The positive correlation between f_{FIR}/f_B and L_B seen in the late-type galaxies may be due to the metallicity effect that less luminous galaxies tend to be metal-poorer (e.g., Roberts, Haynes 1994). The $0 \leq T \leq 5$ group has few sample with $L_B < 10^9 L_\odot$ and the effective span of L_B is nearly half of that for the $6 \leq T \leq 11$ group. We need more data of less luminous galaxies in the $0 \leq T \leq 5$ group in order to examine whether the correlation is a common feature in both the early- and late-type spiral galaxies.

We make an universal f_{FIR}/f_B distribution using the data shown in figure 6. For more distant galaxies, only those having higher f_{FIR}/f_B values are detectable in the *IRAS* survey; we have to correct this ‘‘Malmquist’’ bias. All of our sample galaxies are nearby

galaxies (see table 2) and it is natural to assume no evolution of the $f_{\text{FIR}}/f_{\text{B}}$ frequency distribution with distance among our sample. Then, the $f_{\text{FIR}}/f_{\text{B}}$ frequency distribution of objects in a small range of B magnitude represents the universal $f_{\text{FIR}}/f_{\text{B}}$ distribution of galaxies in the Local Universe. We make $f_{\text{FIR}}/f_{\text{B}}$ distribution histograms in seven B_{T}^0 -magnitude ranges, i.e., 9 - 10 mag, 10 - 11 mag, 11 - 12 mag, 12 - 13 mag, 13 - 14 mag, 14 - 15 mag, and 15 - 16 mag; the lower limits of $\log (f_{\text{FIR}}/f_{\text{B}})$ used for the analysis are -2.2 , -1.8 , -1.4 , -1.0 , -0.6 , -0.2 , and 0.2 , respectively. The final frequency histogram is obtained by combining the magnitude-sliced $f_{\text{FIR}}/f_{\text{B}}$ frequency histograms down to the detection limit with a weight of number of the objects. The functions are shown in figure 8 for each morphological type of $0 \leq T \leq 5$. Figure 8 is the firstly derived universal distribution of the present star formation activity in spiral galaxies. The spirals later than $T = 6$ (Scd) contain the possible metallicity effect (see figure 7), and for these galaxies the indicator of the present star formation activity must be the values of $f_{\text{FIR}}/f_{\text{B}}$ with some correction of this effect depending on L_{B} . But this is a subject in future work.

The extent of $\log (f_{\text{FIR}}/f_{\text{B}})$ is from about -2 to 0.5 for S0a and Sa, from -1.5 to 0.5 for Sab and Sb, and from -1.0 to 0 for Sbc and Sc. The corresponding present-to-past averaged SFRs range from almost zero to ten for S0a and Sa, from a tenth to ten for Sab and Sb, and from several tenths to several for Sbc and Sc. The dispersion of $\log (f_{\text{FIR}}/f_{\text{B}})$ in each morphological type is much larger than the difference between the morphological types. For instance, the median of $\log (f_{\text{FIR}}/f_{\text{B}})$ is -0.6 in Sa, and -0.4 in Sb, and the difference is only 0.2 , while the dispersion is about 2 in both Sa and Sb. The mean value of the $f_{\text{FIR}}/f_{\text{B}}$ is similar to each other between morphological types, which is in contrast with the suggestion by de Jong et al. (1984). It is interesting to note that the dispersion of $(B - V)_{\text{T}}^0$ in each morphology is comparable to the difference of the median value between morphologies and that the dispersion of $(B - V)_{\text{T}}^0$ in each morphology, corresponding to $\Delta \log (f_{\text{B}}/f_{\text{V}}) \sim 0.1$, is much smaller than the dispersion of $\log (f_{\text{FIR}}/f_{\text{B}})$ (compare figure 1 with figure 8). The shape of the distribution histogram of $f_{\text{FIR}}/f_{\text{B}}$ is different

from morphological type to type; the histogram is Gaussian-like for Sbc and Sc samples, while the shape is flat for S0a and Sa samples, indicating the star formation mechanism in S0a and Sa is different from that in Sbc and Sc.

Kennicutt (1983) derived a similar distribution of the present-to-past averaged SFRs for about 100 galaxies, using the $H\alpha$ luminosity, the total mass estimated using B luminosity, and the morphology-dependent mass-to-luminosity ratios (M/L). However, he only pointed out that the present-to-past averaged SFR is higher for later-type spirals. We should emphasize here that the dispersion of the present-to-past averaged SFR has an important meaning that the star formation activity of spiral galaxies, especially early-type spirals, has a large time variation. Coziol and Demers (1995) showed that L_{FIR} has a tight correlation with L_{B} for the starburst galaxies made from Montreal-Blue-Galaxy sample (see their figure 2), and concluded that the starburst galaxies have a nearly constant SFR over Gyr scale. This disagrees with our result. However, their figure must be seen carefully, because the luminosity-luminosity relations tend to have a tight correlation in general. From our analysis, the starburst galaxies have high $f_{\text{FIR}}/f_{\text{B}}$ values i.e., $\log(f_{\text{FIR}}/f_{\text{B}}) \sim 0$ for any L_{B} and morphological types, and thus necessarily shows a tight correlation between L_{FIR} and L_{B} .

4 Discussion

What kind of mechanism causes the large variation of the present star formation activity especially for early-type spirals? In this section, we discuss the correlation between the present star formation activity and the features and environments of galaxies.

4.1 Bar

Whether the bar structure is the effective exciting mechanism for intense star formation is still an open question. In our sample data, 85% of spiral galaxies are divided into three morphological subgroups concerning the bar structure, i.e., SA (non-barred, 21% of the

total sample), SX (mixed, 28%), and SB (barred, 36%). We show in figure 9 the universal frequency histograms of $\log (f_{\text{FIR}}/f_{\text{B}})$ of the barred galaxies (SB sample) for Sa ($T = 1$), Sb ($T = 3$), and Sc ($T = 5$), respectively by solid lines. In each histogram, the dashed lines indicate the distribution of the non-barred galaxies (SA sample). Though the barred galaxy sample has somewhat larger $f_{\text{FIR}}/f_{\text{B}}$ values than the non-barred galaxy sample for the Sa sample, the distributions of the barred and the non-barred samples are similar to each other in all three morphologies. The optical bar structure is, thus, not correlated with the present star formation activity. This disagrees with the result by de Jong et al. (1984), which was derived by using the small number of sample.

4.2 Nuclear and Disk Starburst

It is known that some galaxies show intense starburst in their nuclear regions. Devereux (1989) compiled 132 nearby nuclear-region starburst galaxies according to $10\ \mu\text{m}$ observations. Figure 10 shows the $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ diagrams for the nuclear-starburst sample by Devereux (1989) among our sample for Sa, Sb, and Sc. The dashed symbols indicate the total sample and the bold solid circles the nuclear-region starburst sample among our sample. It is seen that the nuclear-region starburst galaxies tend to have the higher values of $f_{\text{FIR}}/f_{\text{B}}$ in all three morphologies. The existence of the post-nuclear-starburst galaxies (e.g., Walker et al. 1987; Beckman et al. 1991) suggests that turning on and off the nuclear-region starburst often happens. Krügel and Tutukov (1993) showed by an analytic modeling that a possibility of repetitive starburst in the nuclear region in a time scale of 10^{7-8} yr.

Disk star formation also contributes to the total star formation activity. Rice et al. (1988) tabulated FIR-to- B diameter-size ratios for galaxies with large angular sizes; the ratio measures the development of the star-forming disk. Figure 11 shows a correlation between the size ratio and the flux ratio; filled marks are for the earlier-type galaxies with $T \leq 4$ (Sbc) and open marks are for the later-type galaxies with $T \geq 5$ (Sc). The larger the FIR-to- B flux ratio is, the larger the FIR-to- B size ratio is; this trend holds for both

earlier and later spirals. Though the number of the sample is small, the expansion of the starburst disk seems to be one of the dominant features of the high total star formation activity as well as the nuclear-region starburst.

The H I gas content may be an important factor of the star formation. RC3 gives the H I index, $\text{H I index} \equiv -2.5 \log (f_{\text{H I}}/f_{\text{B}}) + \text{constant}$, where $f_{\text{H I}}/f_{\text{B}}$ is the flux ratio of the H I 21 cm line and the B band. Note that the higher value of the H I index indicates being more H I-deficient with respect to the B band flux. RC3 cataloged only reliable H I index values and for 1204 galaxies among our sample (72 %). Figure 12 shows a correlation between $\log (f_{\text{FIR}}/f_{\text{B}})$ and the H I index for the 1204 galaxies. The circles indicate 962 objects with reliable $f_{\text{FIR}}/f_{\text{B}}$, and the crosses indicate 242 objects without reliable $f_{\text{FIR}}/f_{\text{B}}$; true positions would be more left on the diagram. For the 962 objects with reliable $f_{\text{FIR}}/f_{\text{B}}$, the gradient is 0.29 or $\Delta \log (f_{\text{H I}}/f_{\text{B}})/\Delta \log (f_{\text{FIR}}/f_{\text{B}}) = 0.29 / (-2.5) \sim -0.1$ with a correlation coefficient of 0.12; i.e., there is no correlation between $f_{\text{H I}}/f_{\text{B}}$ and $f_{\text{FIR}}/f_{\text{B}}$. This suggests that the time scale of the present star formation activity is much shorter than that of gas consumption in a galaxy of Gyr scale (e.g., Kennicutt et al. 1994).

Kennicutt (1988) and Kennicutt et al. (1989) pointed out that the more intense star-forming galaxies have more numerous, more luminous, and larger H II regions. They suggested that this is due to the variation of the mass function of the molecular clouds. M 51 is an Sbc ($T = 4.0$) galaxy with $\log (f_{\text{FIR}}/f_{\text{B}}) = -0.14$, a high value for Sbc sample (see figures 3 and 8 (e)). Adler et al. (1992) mapped M 51 by ^{12}CO ($J = 1 - 0$) line using an interferometer with a spatial resolution of $7'' \times 11''$ (350×550 pc at a distance of 9.6 Mpc), and found many associations of giant molecular clouds with sizes of 300 - 800 pc and masses of $(1 - 40) \times 10^7 M_{\odot}$, which are one order of magnitude larger as compared with the Galactic one. However, the high resolution mappings by the molecular lines have been made for only a few galaxies so far. High resolution systematic surveys of the molecular gas in many galaxies are necessary to study the degree of the gas concentration

in the central regions and the mass functions of the molecular clouds in the disks.

4.3 Tidal Distortion

The galaxy-galaxy interaction is considered to be a trigger for the central-region starburst from observations (e.g., Heisler, Vader 1995) and from numerical simulations (e.g., Wada, Habe 1992). We examine our sample galaxies appearing in catalogs by Arp (1966) and by Vorontsov-Velyaminov (1977) in which they collected the distorted-shaped galaxies. Figures 13 and 14 show $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ diagrams for the Arp sample and the Vorontsov-Velyaminov sample, respectively, in the same way as figure 10. For Sb galaxies, both the distorted-shaped samples tend to show high $f_{\text{FIR}}/f_{\text{B}}$ values. However, for Sa and Sc galaxies, the distorted-shaped samples show no different distribution from that of the total sample. 60% of the Arp sample shown in figure 13 and 68% of the Vorontsov-Velyaminov sample shown in figure 14 share the common sample. The common sample has also a similar behavior on $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ diagram. Galaxies with distorted shape, thus, does not always indicate the higher present star formation activity.

4.4 Spatial Number Density of Galaxies

The spatial number density of galaxies is considered to affect the star formation history of the galaxies. We investigate the galaxies in the three extreme environments, which are field isolated galaxies, compact group members, and cluster members. We refer to the isolated sample catalog by Karachentseva (1973). Figure 15 shows a $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ diagram in the same way as figure 10. The number of the Karachentseva sample for Sa is only two. For Sb and Sc galaxies, the distribution of the Karachentseva sample is similar to that of the total sample. It seems, therefore, that the environment of “extreme field” does not affect the star formation activity.

Hickson (1982, 1993) made the Compact Group catalog. The compact groups give the most highest spatial number density environment in 10 kpc scale, though the group itself is isolated in Mpc scale. We have omitted from our sample the companion systems that

we can not identify the association of *IRAS* point sources on such individual objects, as mentioned in section 2.1, but some galaxies in relatively loose compact groups are left in our sample. Figure 16 shows a $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ diagram for the Hickson sample in the same way as figure 10. Though the number of the sample is small, it seems that there is no significant difference on this diagram between the group member sample and the total sample.

For the cluster member sample, we refer to the rich cluster catalog by Abell et al. (1989). We pick up cluster members by satisfying both of two criteria that (1) the location is within 0.5 Abell radius ($1 R_{\text{A}} \equiv 1.7z^{-1}$ arcmin, z is the redshift of the cluster) from the center of the cluster, and that (2) the radial velocity difference is within 500 km s^{-1} with respect to the mean cluster radial velocity. This procedure picks up effectively galaxies in the central region of the rich clusters. Since our sample galaxies are mostly nearby objects (see table 2), they contain a small number of Abell cluster members. We add three nearby poor clusters, Virgo, Cancer, and Pegasus I clusters as the cluster sample; the criteria of the membership are the same as for the Abell cluster members. Figure 17 shows a $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ diagram in the same way as figure 10. Though the number of the cluster member is small, again it seems that there is no significant difference on the diagram between the cluster member sample and the total sample. Thus, the spatial number density of galaxies does not correlate with the activity of the present star formation of spiral galaxies.

5 Summary

We constructed a new large data set of 1681 nearby spiral galaxies from available databases, and studied the variation of the present star formation activity. Using these data and the FIR and *B*-band surface intensity image data of M 31, we showed that $f_{\text{FIR}}/f_{\text{B}}$ is a useful indicator of the present star formation activity in spiral galaxies; $\log (f_{\text{FIR}}/f_{\text{B}}) \simeq -2$ corresponds to no present star formation activity, where all FIR

emission is due to the cirrus component, and $f_{\text{FIR}}/f_{\text{B}}$ increases with the present star formation activity. The value of $\log (f_{\text{FIR}}/f_{\text{B}})$ would be about -0.7 to -0.6 in the case that the present SFR is the same as the past-averaged SFR and the maximum value of $\log (f_{\text{FIR}}/f_{\text{B}})$ amounts to about 0.5 . Analyzing the distribution of $f_{\text{FIR}}/f_{\text{B}}$ of the sample galaxies, we showed that the present star formation activity has a great variety even within the same morphological type of galaxies and that the star formation activity changes discontinuously in a short time scale. We also showed that the intense star formation activity does not continue more than 10^8 yr. For spirals with $T = 4$ to 5 , the time variation of the present star formation activity is not so large as that for spirals with $T = 0$ to 1 . We have made a universal frequency histogram of $\log (f_{\text{FIR}}/f_{\text{B}})$, the present star formation activity indicator, for each morphological type of spiral galaxies. We pointed out that the shape of the universal histogram of $\log (f_{\text{FIR}}/f_{\text{B}})$ in Sbc and Sc sample is Gaussian-like, while that in S0a and Sa sample is fairly flat. The systematic difference of the $\log (f_{\text{FIR}}/f_{\text{B}})$ distribution along the morphological type suggests that the mechanisms regulating the present star formation activity are internal processes in galaxies; the different morphological type may cause the different star formation mechanism and the different star formation mechanism may maintain the morphology.

We investigated what structures and environments of galaxies are correlated with the present star formation activity. The nuclear-region starburst may be a state of the intense star formation activity in the whole galaxy. The disk-wide starburst also has an important contribution to the total star formation activity. The environmental effects are little correlated with the present star formation activity, though the galaxy-galaxy interaction is considered to be a triggering mechanism for the intense star formation in galaxies. This is consistent with the suggestion mentioned above that the intense star formation does not continue for a long time.

We would like to thank Keiichi Kodaira for critical reading of the manuscript and Hidemi Suzuki for kindly allowing us to use the B band image of M 31. We made use of

NDADS/ARMS on-line service and thank Walter Rice, Melissa Larkin, and Seth Digel for their kind guide. This research has also made use of the NASA/IPAC Extragalactic Database (NED) which is operated by the Jet Propulsion Laboratory, Caltech, under contact with the National Aeronautics and Space Administration.

Appendix 1. Relation of $\log (f_{\text{FIR}}/f_{\text{B}})$ vs Inclination of Galaxies

The B_{T}^0 -system in RC3 corrects statistically the internal absorption in galaxy depending on apparent inclination of galaxy. We investigate the distribution of $\log (f_{\text{FIR}}/f_{\text{B}})$ as a function of apparent inclination of galaxy to check whether the distribution of $\log (f_{\text{FIR}}/f_{\text{B}})$ is seriously affected by the extinction correction. Apparent inclinations of galaxies versus $\log (f_{\text{FIR}}/f_{\text{B}})$ diagrams are shown for Sa ($T = 1$), Sb ($T = 3$), and Sc ($T = 5$) samples in figure 18 (a) and for the total 1681 sample in figure 18 (b). We obtain the apparent inclination of a galaxy, i [degree], from apparent ratio of minor-to-major axes at a surface brightness of $25 \text{ mag arcsec}^{-1}$ at B band, R , using an equation shown in *Nearby Galaxies Catalog* by Tully (1988), $\cos^2(i - 3^\circ) = (R^2 - 0.2^2)/(1 - 0.2^2)$. It is found from figure 18 that the mean value and the dispersion of $\log (f_{\text{FIR}}/f_{\text{B}})$ are independent of the inclination.

Appendix 2. The Time Scale of the Intense Star Formation Activity

The actively star-forming galaxies have values of $\log (f_{\text{FIR}}/f_{\text{B}})$ as high as 0 to 0.5. If galaxies would maintain such high activities for a long time, B-type and A-type stars are accumulated in a large number; the plots on the $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ diagrams shown in figures 6 and 7 would move to the lower right direction along the line with a coefficient of -1 .

We estimate the evolution of $f_{\text{FIR}}/f_{\text{B}}$ and L_{B} under an assumption of the constant star formation rate at a high level. We assume the IMF of Salpeter (1955), i.e., $n(m)dm \propto m^{-2.35}dm$, where $n(m)dm$ is the number of stars having masses of m to $m + dm$. The mass versus bolometric luminosity relation of stars is $l(m) \propto m^{3.45}$ (Allen 1973). The total luminosity of stars having masses of m to $m + dm$ is, therefore, $n(m)l(m)dm \propto m^{1.10}dm$. The life time of a star is written as $t(m) \propto m/l(m) \propto m^{-2.45}$, or $t(m) = 10^{10}m^{-2.45} \text{ yr}$,

for m in the unit of $M_\odot$. The starburst forms stars with masses from the lowest one of m_l to the most massive one of m_u . Let t_0 be the time since the beginning of the constant starburst, the resultant total luminosity relative to the luminosity by a single-generation starburst, the time of which is put here by a life time of the most massive star, is $L/L_{\text{sing}} \equiv (\int_{m_0}^{m_u} m^{1.10} m^{-2.45} dm + m_0^{-2.45} \int_{m_l}^{m_0} m^{1.10} dm) / (m_u^{-2.45} \int_{m_l}^{m_u} m^{1.10} dm)$, where m_0 is a mass of the most massive star which has survived for the starburst duration, t_0 , and usually $m_u > m_0 \gg m_l$. Note that stars with masses around m_0 contribute the most to the total luminosity. We also assume that the FIR luminosity generated by the recent starburst dominates the observed FIR luminosity because of very intense present star formation activity. By such a continuous intense star formation, the total luminosity L increases with time, and L/L_{sing} would be about ten when $t_0 \sim 10^8$ yr (corresponding to $m_0 \sim 6 M_\odot$) for $m_u = 10$ to $50 M_\odot$. In this time, L nearly corresponds to L_B because of blue colors with $m_0 \sim 6 M_\odot$ and L_{sing} to L_{FIR} ; i.e., L_B has increased by a factor of ten and f_{FIR}/f_B has decreased by a factor of ten. There would be, therefore, a “forbidden” region in the upper right corner of the $\log(f_{\text{FIR}}/f_B)$ versus $\log L_B$ distribution of galaxies. This is not the case in figures 6 and 7 and thus the intense star formation activity of a galaxy continues only shorter than 10^8 yr.

References

- Abell G.O., Corwin H.G., Jr., Olowin R.P. 1989, ApJS, 70, 1
- ADC CD-ROM Selected Astronomical Catalogs Volume 1* prepared by L.E. Brotaman, S.E. Gessner, NASA/GSFC Astronomical Data Center
- Adler D.S., Lo K.Y., Wright M.C.H., Rydbeck G., Plante R.L., Allen R.J. 1992, ApJ, 392, 497
- Allen C.W. 1973, *Astrophysical Quantities Third Edition* (The Athlone Press, University of London, London)
- Arimoto N., Yoshii Y. 1986, A&A, 164, 260
- Arp H.C. 1966, ApJS, 14, 1
- Beckman J.E., Varela A.M., Muñoz-Tuñon C., Vílchez J.M., Cepa J. 1991, A&A, 245, 436
- Beichman C.A. 1987, ARA&A, 25, 521
- Beichman C.A., Myers P.C., Emerson J.P., Harris S., Mathieu R., Benson P.J., Jennings R.E. 1986, ApJ, 307, 337
- Belfort P., Mochkovitch R., Dennefeld M. 1987, A&A, 176, 1
- Bothun G.D., Lonsdale C.J., Rice W. 1989, ApJ, 341, 129
- Bruzual A.G. 1983, ApJ, 273, 105
- Campos-Aguilar A., Moles M. 1991, A&A, 241, 358
- Cataloged Galaxies and Quasars Observed in the IRAS Survey* 1985, prepared by C.J. Lonsdale, G. Helou, J. Good, W. Rice (JPL, Pasadena) (CGQIRAS)
- Coziol R., Demers S. 1995, ApL&Comm, 31, 41
- de Jong T., Clegg P.E., Soifer B.T., Rowan-Robinson M., Habing H.J., Houck J.R., Aumann H.H., Raimond E. 1984, ApJL, 278, L67
- Devereux N.A. 1989, ApJ, 346, 126
- Devereux N.A., Jacoby G., Ciardullo R. 1995, AJ, 110, 1115
- Devereux N.A., Price R., Wells L.A., Duric N. 1994, AJ, 108, 1667

- Devereux N.A., Scowen P.A. 1994, AJ, 108, 1244
- Devereux N.A., Young J.S. 1990, ApJL, 350, L25
- Devereux N.A., Young J.S. 1991, ApJ, 371, 515
- Dressler A., Thompson I.B., Shectman S.A. 1985, ApJ, 288, 481
- Fabbiano G., Gioia I.M., Trinchieri G. 1988, ApJ, 324, 749
- Gallagher J.S., III, Hunter D.A., Tutukov A.V. 1984, ApJ, 284, 544
- Gavazzi G., Boselli A., Kennicutt R. 1991, AJ, 101, 1207
- Gavazzi G., Jaffe W. 1985, ApJL, 294, L89
- Guiderdoni G., Rocca-Volmerange B. 1987, A&A, 186, 1
- Hamabe M., Ichikawa S. 1992, in Proc of Astronomical Data Analysis Software and System I, ASPC, 25, ed D. Worrall et al. (ASP, San Francisco) p325
- Heisler C.A., Vader J.P. 1995, AJ, 110, 87
- Hickson P. 1982, ApJ, 255, 382
- Hickson P. 1993, ApL&Comm, 29, 1
- IRAS Faint Source Catalog* $|b| > 10$ Degrees Version 2.0 1989, prepared by M. Moshir (GPO, Washington, DC) (FSC)
- IRAS Point Source Catalog* 1985, Joint *IRAS* Science Working Group (GPO, Washington, DC) (PSC)
- IRAS Small Scale Structures Catalog* 1986, prepared by G. Helou, D.W. Walker (GPO, Washington, DC) (SSSC)
- Karachentseva V.E. 1973, Soob. Spets. Astrofiz. Obs. (Communication of the Spetial Astrophysical Observatory), No.8
- Keel W.C. 1993, AJ, 106, 1771
- Kenney J.D.P., Wilson C.D., Scoville N.Z., Devereux N.A., Young J.S. 1992, ApJL, 395, L79
- Kennicutt R.C., Jr. 1988, ApJ, 334, 144
- Kennicutt R.C., Jr. 1983, ApJ, 272, 54

- Kennicutt R.C., Jr. 1992, ApJ, 388, 310
- Kennicutt R.C., Jr., Edgar B.K., Hodge P.W. 1989, ApJ, 337, 761
- Kennicutt R.C., Jr., Kent S.M. 1983, AJ, 88, 1094
- Kennicutt R.C., Jr., Tamblyn P., Congdon C.W. 1994, ApJ, 435, 22
- Krügel E., Tutukov A.V. 1993, A&A, 275, 416
- Larson R.B., Tinsley B.M. 1978, ApJ, 219, 46
- Moss C., Whittle M. 1993, ApJL, 407, L17
- NAO/ADAC CD-ROM Catalog Volume 1. Galaxies and Non-stellar Objects* 1994, prepared by M. Hamabe, S. Ichikawa, S. Nishimura
- Nearby Galaxies Catalog* 1988, Tully R.B. (Cambridge University Press, Cambridge)
- Persson C.J.L., Helou G. 1987, ApJ, 314, 513
- Rice W. 1993, AJ, 105, 67 (*IRAS HiRes*)
- Rice W., Lonsdale C.J., Soifer B.T., Neugebauer G., Kopan E.L., Lloyd L.A., de Jong T., Habing H.J. 1988, ApJS, 68, 91
- Roberts M.S., Haynes M.P. 1994, ARA&A, 32, 115
- Roberts W.W., Jr., Huntley J.M., van Albada G.D. 1979, ApJ, 233, 67
- Salpeter E.E. 1955, ApJ, 121, 161
- Sandage A. 1986, A&A, 161, 89
- Sauvage M., Thuan T.X. 1992, ApJL, 396, L69
- Sauvage M., Thuan T.X. 1994, ApJ, 429, 153
- Searle L., Sargent W.L.W., Baguolo W.G. 1973, ApJ, 179, 427
- Shu F.H., Adams F.C., Lizano S. 1987, ARA&A, 25, 23
- Soifer B.T., Boehmer L., Neugebauer G., Sanders D.B. 1989, AJ, 98, 766
- Soifer B.T., Sanders D.B., Madore B.F., Neugebauer G., Danielson G.E., Elias J.H., Lonsdale C.J., Rice W.L. 1987, ApJ, 320, 238
- Third Reference Catalogue of Bright Galaxies* 1991, de Vaucouleurs G., de Vaucouleurs A., Corwin H.G., Jr., Buta R.J., Paturel G., Forqué P. (Springer-Verlag, New York)

- Vorontsov-Velyaminov B.A. 1977, A&AS, 28, 1
- Wada K., Habe A. 1992, MNRAS, 258, 82
- Walker C.E., Lebofsky M.J., Rieke G.H. 1988, ApJ, 325, 687
- Wood D.O.S., Churchwell E. 1989, ApJ, 340, 265
- Xu C. 1990, ApJL, 365, L47

Figure Captions

Figure 1. Frequency histogram of $(B - V)_T^0$ of galaxies for each morphological type.

The morphological type and index are shown at the right of each diagram. The later the morphological type is, the bluer the color is. The dispersion of the color in each morphological type is small, corresponding to $\Delta \log (f_B/f_V) \sim 0.1$.

Figure 2. A $\log (f_{100}/f_{60})$ versus $\log (f_{60}/f_B)$ diagram for 1150 sample galaxies with reliable f_{60} and f_{100} values. A straight line is a regression line and the correlation coefficient is 0.72. Most of the objects showing the large deviation from the regression line are irregular galaxies. Using only objects with $0 \geq T \geq 5$, the correlation coefficient is 0.80. Using the relation of the regression line, $\log (f_{100}/f_{60}) = -0.23 \log (f_{60} [\text{Jy}] / f_B [\text{W m}^{-2}]) + 3.41$, we estimate f_{100} for 156 objects with upper-limited f_{100} and reliable f_{60} .

Figure 3. A $\log (f_{100}/f_{60})$ versus $\log (f_{\text{FIR}}/f_B)$ diagram for 1150 sample galaxies. Large circles indicate 916 early-type spirals ($0 \geq T \geq 5$) and small circles indicate 473 late-type spirals ($6 \geq T \geq 11$). Six famous spiral galaxies without strong AGN are denoted by large filled squares with numbers; 1: M 31 ($T = 3$), 2: M 81 ($T = 2$), 3: M 33 ($T = 6$), 4: M 101 ($T = 6$), 5: M 51 ($T = 4$), 6: M 83 ($T = 5$). The three areas enclosed by solid lines show the color-color relation corresponding to (A) the H II regions, (B) the non-H II regions, and (C) the nuclear regions in M 31, as shown in figure 5. The composition of the H II regions (A) and the non-H II regions (B) in various fraction makes the sequence shown by the bold solid line (see text), which is similar to the sequence of the galaxy distribution on this diagram.

Figure 4. (a) The B -band image taken at Ouda Station, Kyoto University. Foreground stars are removed using IRAF task of ‘imedit’. The north is top and the east is left. The image size is $1.'68 \times 2.'00$. (b) The f_{FIR}/f_B image made by the B -band image shown in (a) and *IRAS HiRes* image data of Rice (1993). The sample locations for

measuring the surface intensities in FIR and B bands are indicated by circles. The size of the circles corresponds to the aperture size for the photometry, $0.''5$ in radius. The numbers 1 - 15 represent H II regions, 16 - 25 non-H II regions in disk, 26 - 32 bulge regions, and 33 - 35 central regions.

Figure 5. The same as figure 3 but for various areas in M 31 with an aperture radius of $0.''5$, i.e., 15 points from H II regions in disk (2 to 8 kpc from the nucleus), 10 points from non-H II regions in disk (2 to 8 kpc), 7 points from bulge region (1 to 2 kpc), and 3 points from nuclear region (within 0.3 kpc), the locations of which are shown in figure 4. The meanings of the symbols are shown in the diagram. The solid line (A) encloses the H II regions, the solid line (B) encloses both the non-H II regions in disk and bulge regions, which occupy the same area and are both called as non-H II regions. The solid line (C) encloses the three points in the central 0.3 kpc. The big star mark is the position of the whole M 31, which is inside the region B, indicating that most of the FIR emission of M 31 comes from cirrus component.

Figure 6. A $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ diagram for each morphological type, from (a) $T = 0$ (S0a) to (k) $T = 10, 11$ (Im/cI). Circles indicate objects with measured f_{FIR} , while crosses indicate ones with upper-limited f_{FIR} . Size of the symbols shows B_{T}^0 magnitude as shown in the upper left corner of the diagram. Note that L_{B} is written in the unit of solar luminosity; $L_{\odot} = 3.85 \times 10^{26}$ W. Small tick marks near the left ordinate show the detection limit of $f_{\text{FIR}}/f_{\text{B}}$ due to the *IRAS* detection limit, for given apparent B_{T}^0 magnitude, $B_{\text{T}}^0 = 16, 15, 14, 13, 12, 11, 10$, and 9 mag, respectively from upper to lower.

Figure 7. The same as figure 6 but for the $0 \leq T \leq 5$ group and the $6 \leq T \leq 11$ group. For the early-type spirals, $f_{\text{FIR}}/f_{\text{B}}$ distributes without L_{B} -dependence, while for the late-type spirals, $f_{\text{FIR}}/f_{\text{B}}$ tends to increase with increasing L_{B} . Most of the upper limit sample in the $6 \leq T \leq 11$ group are irregulars; see figure 6 (k).

Figure 8. A universal frequency histogram of $\log (f_{\text{FIR}}/f_{\text{B}})$ for each morphological type.

From upper to lower panels, (a) $T = 0$ (S0a), (b) $T = 1$ (Sa), (c) $T = 2$ (Sab), (d) $T = 3$ (Sb), (e) $T = 4$ (Sbc), (f) $T = 5$ (Sc). We show only objects earlier than Sc, because for the later-type galaxies, $f_{\text{FIR}}/f_{\text{B}}$ depends on L_{B} . The dashed line indicates the upper limit value. $\log (f_{\text{FIR}}/f_{\text{B}}) \simeq -2$ corresponds to no present star formation activity. The histograms are Gaussian-like for Sbc and Sc samples, while they are rather flat for S0a and Sa samples.

Figure 9. A universal frequency histogram of $\log (f_{\text{FIR}}/f_{\text{B}})$ for the barred (SB) sample (solid line) and the non-barred (SA) sample (dashed line). Only Sa ($T = 1$), Sb ($T = 3$), and Sc ($T = 5$) samples are shown here. The number of the barred sample is 65, 117, and 78 for $T = 1, 3$, and 5 , respectively, and the number of the non-barred sample is 38, 55, and 84, respectively. There is no significant difference between both samples.

Figure 10. A $\log (f_{\text{FIR}}/f_{\text{B}})$ versus $\log L_{\text{B}}$ diagrams for the nuclear-region starburst galaxies listed in Devereux (1989). Size of the symbols shows B_{T}^0 magnitude as shown in the upper left corner of the diagram and the magnitude ranges are the same as those in figure 6. The bold solid symbols indicate the nuclear-region starburst sample. The dashed symbols indicate the total sample, the same as shown in figure 6. Generally the nuclear-starburst sample has high $f_{\text{FIR}}/f_{\text{B}}$ values.

Figure 11. A $\log (f_{\text{FIR}}/f_{\text{B}})$ versus FIR-to-B size ratio diagram for the galaxies listed in Rice et al. (1988). Filled circles are for the earlier-type galaxies with $T \leq 4$ (Sbc) and open circles are for the later-type galaxies with $T \geq 5$ (Sc). The more the FIR-disk develops, the larger the star formation indicator, $f_{\text{FIR}}/f_{\text{B}}$, is.

Figure 12. A $\log (f_{\text{FIR}}/f_{\text{B}})$ versus the H I index diagram for 1204 galaxies with the H I index values cataloged in RC3. The circles indicate the objects with reliable $f_{\text{FIR}}/f_{\text{B}}$, and the crosses indicate the objects without reliable $f_{\text{FIR}}/f_{\text{B}}$; true positions would be

more left on the diagram. There is no correlation between $\log (f_{\text{FIR}}/f_{\text{B}})$ and the H I index.

Figure 13. The same as figure 10 but for the peculiar-shaped galaxies listed in Arp (1966). In Sb panel, the Arp sample shows a high $f_{\text{FIR}}/f_{\text{B}}$ value compared with the total sample, however, in Sa and Sc panels, the Arp sample does not show any peculiar distribution on the diagrams.

Figure 14. The same as figure 10 but for the peculiar-shaped galaxies listed in Vorontsov-Velyaminov (1977). The trend is almost the same as that for the Arp sample in figure 13.

Figure 15. The same as figure 10 but for the field isolated galaxies listed in Karachentseva (1973). The number of the Karachentseva sample is only two for Sa. For Sb and Sc, there is no significant difference of distribution on the diagram between the Karachentseva and the total samples.

Figure 16. The same as figure 10 but for the compact group members listed in Hickson (1982, 1993). Though the number of group members is small, it seems that they do not have different distribution from that of the total sample on the diagram.

Figure 17. The same as figure 10 but for the cluster members. We take into account rich clusters cataloged by Abell et al. (1989) and three poor clusters, Virgo, Pegasus I, and Cancer. Though the number of cluster members are small, it seems that they do not have different distribution from that of the total sample.

Figure 18. Distributions of $\log (f_{\text{FIR}}/f_{\text{B}})$ as a function of apparent inclinations of galaxies (a) for Sa ($T = 1$), Sb ($T = 3$), and Sc ($T = 5$) samples and (b) for the total 1681 sample. Open circles are for galaxies with reliable $f_{\text{FIR}}/f_{\text{B}}$ and crosses indicate the upper limits of $f_{\text{FIR}}/f_{\text{B}}$. Inclinations of 0° and 90° mean face-on and edge-on, respectively.

Table 1. The morphology and magnitude distribution of the sample.

T^*	$B_T^{0\dagger}$								Total
	- 9	9 - 10	10 - 11	11 - 12	12 - 13	13 - 14	14 - 15	15 -	
0	0	1	6	11	40	25	13	5	101
1	1	1	4	25	45	52	15	9	152
2	3	3	12	20	43	24	15	1	121
3	1	13	21	43	77	70	24	13	262
4	2	4	25	60	72	67	20	7	257
5	1	4	27	58	103	94	24	4	315
6	4	5	11	32	29	27	18	3	129
7	3	5	3	15	13	10	11	1	61
8	0	1	0	7	15	12	14	1	50
9	3	0	3	10	22	22	27	3	90
10	0	2	5	8	20	41	55	11	142
11 [§]	0	0	0	0	0	0	1	0	1
Total	18	39	117	289	479	444	237	58	1681

* The morphological index given in RC3; i.e., 0:S0a, 1:Sa, 2:Sab, 3:Sb, 4:Sbc, 5:Sc, 6:Scd, 7:Sd, 8:Sdm, 9:Sm, 10:Im, 11:cI.

[†] The classes of B_T^0 magnitude range; - 9 means $B_T^0 < 9.0$, 9 - 10 means $9.0 \leq B_T^0 < 10.0$, and 15 - means $B_T^0 \geq 15.0$.

[§] One object of $T = 11$ is combined into the $T = 10$ group in all analyses in this paper.

Table 2. The morphology and radial velocity distribution of the sample.

T^*	$V_{\text{GSR}} \text{ [km s}^{-1}\text{]}^\dagger$									Total
	< 1000	1000	2000	3000	4000	5000	6000	7000	8000 -	
0	10	32	18	7	10	11	6	3	4	101
1	16	39	22	12	20	21	6	3	13	152
2	15	29	12	19	15	14	4	6	7	121
3	20	51	45	32	38	25	17	3	31	262
4	19	57	43	30	25	25	18	11	29	257
5	29	66	59	24	47	42	13	12	23	315
6	32	29	19	12	12	11	3	3	8	129
7	25	20	5	3	6	0	1	1	0	61
8	11	20	6	6	5	0	1	1	0	50
9	37	36	13	2	1	1	0	0	0	90
10	78	34	15	9	3	0	0	0	2	142
11 [§]	0	0	0	0	0	0	0	0	1	1
Total	292	413	257	156	182	150	70	43	118	1681

* The morphological index given in RC3; i.e., 0:S0a, 1:Sa, 2:Sab, 3:Sb, 4:Sbc, 5:Sc, 6:Scd, 7:Sd, 8:Sdm, 9:Sm, 10:Im, 11:cI.

[†] The radial velocity V_{GSR} [km s⁻¹] system given in RC3 is adopted. < 1000 means $V_{\text{GSR}} < 1000$, 1000 means $1000 \leq V_{\text{GSR}} < 2000$, and 8000 - means $V_{\text{GSR}} \geq 8000$.

[§] One object of $T = 11$ is combined into the $T = 10$ group in all analyses in this paper.